

AD-A166 516

LASER SPECTROSCOPY OF OZONE(U) MASSACHUSETTS INST OF
TECH CAMBRIDGE DEPT OF CHEMISTRY J L KINSEY 18 DEC 85
AFGL-TR-86-0006 F19628-88-K-0088


1/1

UNCLASSIFIED

F/G 7/4

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A166 516

12

AFGL-TR-86-0006

LASER SPECTROSCOPY OF OZONE

James L. Kinsey

Massachusetts Institute of Technology
Department of Chemistry
Cambridge, MA 02139

18 December 1985

Final Report
1 July 1980 - 30 September 1985

DTIC
ELECTE
APR 15 1986
S D
P D


APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

DTIC FILE COPY

AIR FORCE GEOPHYSICS LABORATORY
AIR FORCE SYSTEMS COMMAND
UNITED STATES AIR FORCE
HANSCOM AIR FORCE BASE, MASSACHUSETTS 01731

109

"This technical report has been reviewed and is approved for publication"




DANIEL H. KATAYAMA
Contract Manager
UV Surveillance & Remote Sensing Branch



LEON J. HEROUX
Chief, UV Surveillance & Remote
Sensing Branch

FOR THE COMMANDER



ROBERT A. SKRIVANEK
Director, Ionospheric Physics Division

This report has been reviewed by the ESD Public Affairs Office (PA) and is releasable to the National Technical Information Service (NTIS).

Qualified requestors may obtain additional copies from the Defense Technical Information Center. All others should apply to the National Technical Information Service.

If your address has changed, or if you wish to be removed from the mailing list, or if the addressee is no longer employed by your organization, please notify AFGL/DAA, Hanscom AFB, MA 01731. This will assist us in maintaining a current mailing list.

Do not return copies of this report unless contractual obligations or notices on a specific document requires that it be returned.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

ADA 166516

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE				
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFGL-TR-86-0006	
6a. NAME OF PERFORMING ORGANIZATION Massachusetts Institute of Technology		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Air Force Geophysics Laboratory	
6c. ADDRESS (City, State and ZIP Code) Dept. of Chemistry Cambridge, MA. 02139			7b. ADDRESS (City, State and ZIP Code) Hanscom AFB Massachusetts 01731-5000	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Air Force Geophysics Laboratory		8b. OFFICE SYMBOL (If applicable) LIU	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F19628-80-K-0088	
8c. ADDRESS (City, State and ZIP Code) Hanscom AFB Massachusetts 01731			10. SOURCE OF FUNDING NOS.	
11. TITLE (Include Security Classification) Laser Spectroscopy of Ozone			PROGRAM ELEMENT NO. 61102F	PROJECT NO. 2303
			TASK NO. G1	WORK UNIT NO. AE
12. PERSONAL AUTHOR(S) James L. Kinsey				
13a. TYPE OF REPORT Final		13b. TIME COVERED FROM 1 Jul 80 to 30 Sep		14. DATE OF REPORT (Yr., Mo., Day) 1985 December 18
				15. PAGE COUNT 14
16. SUPPLEMENTARY NOTATION 85				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD 07	GROUP 04	SUB. GR.	Ozone, spectroscopy, UV, fluorescence spectrum, Huggins bands, Hartley continuum.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Data have been obtained for the first time on the fluorescence spectrum of ozone in the Hartley continuum region. This type of data provides information for a new kind of spectroscopy with hitherto unprecedented capabilities for studying both ground state vibrational structure and excited state dissociation dynamics. From this data and theoretical models, qualitative generalizations are made about the potential energy surface of the upper electronic state of the Hartley transition. Experiments have also been conducted to obtain the high resolution spectrum of some of the Huggins bands for jet-cooled ozone. An analysis of the partially resolved rotational structure of these bands indicates that the symmetry of the Huggins bands is the same as that for the Hartley continuum.				
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL Daniel H. Katayama			22b. TELEPHONE NUMBER (Include Area Code) (617)861-4042	22c. OFFICE SYMBOL LIU

TABLE OF CONTENTS

Introduction	1
Personnel Associated With Work	6
Publications	6
Technical Presentations	7

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	



SUMMARY OF RESEARCH

This contract covered the period July 1, 1980 - September 30, 1985. The aim of the work was to improve understanding of the UV and visible spectrum of the atmospheric species O_3 and to improve knowledge of its vibrational levels in the molecule's ground electronic state. Ozone photodissociates readily on absorption of light at any wavelength longer than ~ 850 nm. Consequently, the absorption spectrum is entirely diffuse. There are three principal spectroscopic regions = The Chappuis band (${}^1B_1 \rightarrow \tilde{X}{}^1A_1$, $\sim \lambda 450-850$ nm), the Huggins band (${}^1B_2 \rightarrow \tilde{X}{}^1A_1$, $\sim \lambda 300-360$ nm) and the Hartley band (${}^1B_2 \rightarrow \tilde{X}{}^1A_1$, $\sim \lambda 200-300$ nm). The dominant and by far the most intense of these is the Hartley band, centered around $\sim \lambda 255$ nm. Because of the rapid predissociation in this region, the band is almost entirely featureless, though there is some very faint structure near the band center. To the red, the much weaker (by $\sim 10^{-3}$) Huggins system is also diffuse, but it exhibits definite vibrational structure.

The original idea of this work was to investigate the feeble structure in the Hartley band and the vibrational transitions in the Huggins band by looking at fluorescence excitation spectra of very cold ozone. The cooling (to ~ 10 K) was to be accomplished by adiabatic expansion of an ozone-helium mixture into a free-jet pulsed molecular beam. Our group already had some experience with O_3 molecular beams and tuneable UV laser spectroscopy.

Introduction:

A summary of the research done for this contract is reported in this technical report. The detailed research results and the description of the experiments have been reported in the open literature and a listing of the journal articles is given in the Publication section of this report.

It is important to note that in the process of conducting laser experiments on ozone, fluorescence was unexpectedly observed when the laser was tuned to 266 nm which is in the peak absorption region of the Hartley continuum. This is surprising because it has always been assumed that the ozone is completely photodissociated in this wavelength region. Apparently, the laser light intensity is sufficiently strong that the small fraction of molecules that fluoresce is observable. This observation has led to a new kind of spectroscopy with unprecedented capabilities for studying excited state dissociation dynamics and ground state vibrational structure.

Shortly after the project started, these goals were expanded to include study of the resolved fluorescence of the photodissociating molecules. Inspired by theoretical ideas put forward by E. J. Heller and coworkers, we realized that such data could provide a new kind of spectroscopy with hitherto unprecedented capabilities for studying both ground-state vibrational structure and excited-state dissociation dynamics. The first measurements of this kind were made on ozone (in a static cell), and they revealed a sequence of vibrational levels in the ground electronic state reaching to within 500 cm^{-1} of the molecule's dissociation limit. This was and remains a record for any polyatomic molecules.

The levels that could be assigned were all of the type $1_n 2_0 3_m$, i.e., no ground-state excited bending levels were observed. In the light of the Heller theory, this is a clear indication that dissociation on the upper potential energy surface proceeds with motion of the departing oxygen atom almost directly in the direction of the broken bond. Further qualitative generalizations were made about the shape of the excited-state potential in the Franck-Condon region. Specifically, it was concluded that the surface is saddle-shaped with respect to q_1 (symmetric stretch) and q_3 (antisymmetric stretch). This confirmed the idea that the weak structure in the Hartley band probably occurs because there are a few oscillations in the q_1 coordinate before the motion accelerates in the q_3 direction leading to dissociation. A more quantitative theoretical treatment has not yet been completed, but such a study is planned for the future. Later work also revealed a very small intensity into the 2_1 level ($\sim 1\%$ of that into 1_1). The quantitative remeasurement of the whole O_3 emission spectrum, with special attention to

such minor features, will be an important step in the continuation of our studies on ozone.

A theoretical model for the stretching vibrations of ozone, based on a completely algebraic Hamiltonian, was completed in collaboration with Professor R. D. Levine. It was found that the algebraic approach could fit the observed levels competitively with conventional (Darling-Dennison) methods, yet retain a consistent anharmonic-oscillator perspective. Darling-Dennison analysis mixes harmonic and anharmonic expressions.

Subsequent experimental work turned more towards the original idea, but focussed on the weak Huggins system instead of the Hartley band. A pulsed nozzle beam appropriate for these studies was developed and tested. At about this time, we also started considering whether some sort of optical multichannel analyzer could be found so that we could record an entire wavelength region simultaneously rather than by tedious scanning from line to line. Because of the extremely low light levels inherent in these experiments, it quickly became clear that the readout error would swamp the signal without at least two stages of amplification (image enhancement). Eventually, a unit that seemed suitable was identified and ordered (with funds from another source). Almost a year expired to delivery of the device, and it was found to have serious design problems as well as manufacturing defects. After a great deal of back and forth with the supplier, this instrument may now be just about ready to test for ozone work.

Meanwhile, the old technique was pursued in our investigation of the Huggins system. When excitation spectra were taken of jet-cooled ozone (the 501 and 600 vibrational bands), the widths of the features narrowed by a

factor of ~ 20 from the room-temperature spectra. This confirmed what we had suspected, that a large part of the room-temperature spectral width was due to rotational congestion. The new linewidth observation allowed a fresh estimate of the predissociative lifetime in the electronically excited state (~ 1 ps).

Further frequency narrowing of the excitation light actually permitted the partial resolution of rotational substructure in both vibrational bands, with individual rotational transitions estimated to be $\sim 1.4 \text{ cm}^{-1}$ wide. This increased the estimate of dissociative lifetime to ~ 4 ps. Because of incomplete thermal equilibration of the jet beam and the intrinsic width of individual rotational lines, these lines could not be fully resolved. Instead, the partially resolved structure was modelled with an asymmetric rotor band contour program. This showed quite clearly that both the $1_0^{5,1}$ and 1_0^6 vibrational transitions have the same rotational structure, which corresponds to an a-axis transition moment. If the equilibrium structure of the electronically excited state were of C_{2v} geometry, spin statistics would make this impossible: the existing rotational states would be a different set for the $1_0^{5,1}$ level than for the 1_0^6 level. Hence, we conclude that the electronically excited state has an equilibrium configuration only of C_s geometry, i.e., the two O-O bonds are of unequal length. In this case, for a high "inversion" barrier, the V_3 quantum numbers would refer to single-well (left or right bond longer) levels. Each level, however, would be split into a closely spaced tunnelling doublet and one of the pair at any V_3 would support the correct rotational states for a-type transitions. Since ab initio calculations indicate a C_s geometry for the $1B_2$ excited state (and not for the $1A_1$ state), we conclude that $1B_2 \tilde{\leftarrow} 1A_1$ is the correct electronic assignment for the

Huggins system. I.e., the Huggins band is actually just the wing of the Hartley band, the former corresponding to transitions into bound regions of the potential energy surface and the latter to transitions into a Franck-Condon region with no excited-state quasibound levels. The details of the rotational band-contour analysis permitted estimates of 1B_2 rotational constants: $A \approx 2.1 \pm 0.04 \text{ cm}^{-1}$, $\bar{B} \approx 0.45.01 \text{ cm}^{-1}$. Without data on isotopic variants these are not sufficient to establish a complete structure, but limits can be set on possible geometrics.

Personnel Associated With Work Performed Under
Contract F19628-80-K-0088

Principal Investigator

James L. Kinsey

Current Position

Professor of Chemistry, M.I.T.

Graduate Students

Dan Imre

Asst. Professor, U. of Washington

Amitabha Sinha

Postdoctoral Fellow, NOAA,
Boulder, CO.

Postdoctoral Associates

James H. Goble

Talandic Research, Pasadena, CA

Michael O. Hale

Postdoctoral Associate, M.I.T.

Publications:

1. "Spectroscopic Characterization of Repulsive Potential Energy Surfaces: Fluorescence Spectrum of Ozone"
D. G. Imre, J. L. Kinsey, R. W. Field, D. H. Katayama
J. Phys. Chem. 86, 2564 (1982).
2. "Chemical Dynamics Studied by Emission Spectroscopy of Dissociating Molecules"
D. Imre, J. L. Kinsey, A. Sinha and J. Krenos
J. Phys. Chem. 88, 3956 (1984).
3. "Excitation Spectroscopy of Jet-Cooled Ozone: The Huggins System"
A. Sinha, D. Imre, J. H. Goble and J. L. Kinsey
to be submitted to Journal of Chemical Physics.

Technical Presentations on Work Sponsored by Contract F19628-80-K-0088

Dam Imre: ACS Annual Meeting 1985
Gordon Conference on Molecular Electronic Spectroscopy 1985
University of Washington 1984
Princeton University 1984
University of Texas 1984
University of Pittsburgh 1984
University of Oregon 1984

Amitabha Sinha: M.I.T. 1985
University of Colorado 1985
Air Force Geophysics Laboratory Ozone Meeting 1983

James L. Kinsey: University of Indiana - Gucker Memorial Lecture 1985
University of Illinois 1985
UCLA/USC/Caltech joint seminar 1985
University of Toronto 1985
University of Pennsylvania 1984
Harvard University 1984
Rutgers University 1984
XVI Informal Photochemistry Conference 1984
University of Rochester 1984
Dartmouth College 1984
Texas A&M University 1983
Rice University 1983
Tulane University 1983
University of California, Berkeley 1983
Stanford University 1983
University of Colorado 1983
Colby College 1983

END

DTic

5-86